



**PRESS RELEASE (2024/09/09)**

## **Most detailed study yet of seismic activity links fault strength to likelihood of large earthquakes**

Seismic observations from over 1000 locations have revealed a link between fault strength and earthquake magnitude, and provide a hint to detecting how close a region is to experiencing a large earthquake.

Fukuoka, Japan – Located on the Pacific Ring of Fire, Japan is one of the most earthquake-prone countries in the world, with over 100,000 small earthquakes occurring each year, and the continuous threat of a “big one”. Currently, predicting when major earthquakes will occur isn’t possible, but by studying the numerous small earthquakes that occur, seismologists in Japan hope to understand more about the processes in the Earth’s crust that lead to major quakes.

Now, researchers from Kyushu University and the University of Tokyo, Japan, have studied seismic activity at an unprecedented level of detail, identifying a link between fault strength and earthquake magnitude. Published in [\*Nature Communications\*](#), the study proposes that the strength of the fault affects the *b*-value — and therefore the likelihood of a major earthquake.

“The *b*-value is a very important constant in seismology that characterizes the relationship between earthquake frequency and size,” explains Professor [Satoshi Matsumoto](#), first author of the study and the Director of Kyushu University’s [Institute of Seismology and Volcanology](#). “If there is a low *b*-value, this means there is a higher proportion of large earthquakes, while a high *b*-value means there is a higher proportion of smaller earthquakes.”

The *b*-value can vary between different locations and also over time, and is often reported to decrease just before a major earthquake. A previous study suggested that the decrease in *b*-value was caused by the increasing stress forces exerted on the fault. Now, this study suggests that fault strength is also a contributing factor.

In the study, the research teams analyzed the seismic action happening in the area around the epicenter of the Western Tottori Earthquake, which occurred in 2000 with a magnitude of 7.3. By installing more than 1000 seismic stations in the area, the researchers could conduct seismic observations with an unprecedented level of accuracy.

“Even two decades on, hundreds of tiny aftershocks still occur, most too small for us to feel,” says Matsumoto.

With so many sensors, the researchers could detect tiny movements of the faults, and also the orientation of each fault within the Earth’s crust.

Using this multitude of data, the team were able to estimate the stress field (the different directions of stress forces exerted on each fault at the time of failure) and allowed them to characterize the faults as strong or weak.

“Under certain stress conditions on each tectonic regime, there is a favorable direction of the

fault plane to slip. When faults are in unfavorable directions, this suggests that these are weak faults that can slip more easily. On the other hand, strong faults require more stress to slip, and have a much more characteristic direction,” explains Matsumoto.

From the stress field calculations, the researchers were also able to estimate the  $b$ -value of event group categorized by fault strength. They found that stronger faults have smaller  $b$ -values, suggesting that large earthquakes are more likely to occur, while weaker faults had larger  $b$ -values, suggesting that major earthquakes are less likely.

“Simply put, these weak faults will likely slip before a large amount of stress builds up, which means that they aren’t able to release a large amount of force,” says Matsumoto.

Through deeper understanding of the factors that impact  $b$ -values, the researchers hope that they will be able to inch closer to the “holy grail” of predicting earthquakes.



“I don’t think we will ever know exactly when an earthquake will strike, but looking at data such as fault direction and fault strength, and calculating  $b$ -values, could help us estimate when a fault has reached a critical point— where just a tiny extra nudge of force is needed for the fault to slip,” concludes Matsumoto. “This information is vital to know in order to be prepared for major earthquakes.”

###

For more information about this research, see "Strength dependency of frequency–magnitude distribution in earthquakes and implications for stress state criticality," Satoshi Matsumoto, Yoshihisa Iio, Shinichi Sakai & Aitaro Kato, *Nature Communications*, [10.1038/s41467-024-49422-7](https://doi.org/10.1038/s41467-024-49422-7)

### **About Kyushu University**

Founded in 1911, [Kyushu University](https://www.kyushu-u.ac.jp/) is one of Japan's leading research-oriented institutes of higher education, consistently ranking as one of the top ten Japanese universities in the Times Higher Education World University Rankings and the QS World Rankings. The university is one of the seven national universities in Japan, located in Fukuoka, on the island of Kyushu—the most southwestern of Japan’s four main islands with a population and land size slightly larger than Belgium. Kyushu U’s multiple campuses—home to around 19,000 students and 8000 faculty and staff—are located around Fukuoka City, a coastal metropolis that is frequently ranked among the world's most livable cities and historically known as Japan's gateway to Asia. Through its [VISION 2030](#), Kyushu U will “drive social change with integrative knowledge.” By fusing the spectrum of knowledge, from the humanities and arts to engineering and medical sciences, Kyushu U will strengthen its research in the key areas of decarbonization, medicine and health, and environment and food, to tackle society’s most pressing issues.



[Contact]

Satoshi Matsumoto, Professor

Faculty of Sciences, Institute of Seismology and Volcanology

Tel: +81-95-762- 6621

E-mail: matumoto@sevo.kyushu-u.ac.jp